

PL-TR-96-2301

CONTROL OF CONTAMINANTS AROUND SPACECRAFT

**Dale J. Levandier
Skip Williams**


**ORION International Technologies, Inc
6501 Americas Parkway NE, Suite 200
Albuquerque, NM 87110**

6 November 1996

19970428 217

Scientific Report No. 1

Approved for public release; distribution unlimited




**PHILLIPS LABORATORY
Directorate of Geophysics
AIR FORCE MATERIEL COMMAND
HANSCOM AFB, MA 01731-3010**

DTIC QUALITY INSPECTED 1

"This technical report has been reviewed and is approved for publication"



RAINER A. DRESSLER
Contract Manager



LEILA S. JEONG
Chief
Simulation Branch



WILLIAM A. M. BLUMBERG
Director
Optical Environment Division

This report has been reviewed by the ESC Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center (DTIC). All others should apply to the National Technical Information Service (NTIS).

If your address has changed, if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify PL/TSB, 29 Randolph Road, Hanscom AFB, MA 01731-3010. This will assist us in maintaining a current mailing list.

Do not report copies of this report unless contractual obligations or notices on a specific document require that it be returned.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 6 November 1997		3. REPORT TYPE AND DATES COVERED Scientific Report No. 1
4. TITLE AND SUBTITLE Control of Contaminants Around Spacecraft			5. FUNDING NUMBERS PE 61102F PR 2303 TA GJ WU AA Contract F19628-92-C-0161	
6. AUTHOR(S) Dale J. Levandier Skip Williams				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) ORION International Technologies, Inc 6501 Americas Parkway NE, Suite 200 Albuquerque, NM 87110			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Phillips Laboratory 29 Randolph Road Hanscom AFB, MA 01731-3010 Contract Manager: Rainer Dressler/GPO			10. SPONSORING/MONITORING AGENCY REPORT NUMBER PL-TR-96-2301	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) A high temperature octopole/collision cell apparatus (HT8P) was developed, along with the required cooling/power feedthroughs and modified "injector lens". The radio frequency operating characteristics of the HT8P were optimized, and the collision cell effective interaction length was determined in a series of calibration experiments. The well known $\text{Ar}^+ + \text{D}^2 \rightarrow \text{ArD}^+ + \text{D}$ reaction was used in the calibration experiments, in which the HT8P was operated at temperatures up to 630 K (675°). The collision cell gas density versus temperature dependence for measurements with a capacitance manometer was modeled theoretically and verified experimentally using the same reaction. This dependence, due to thermal transpiration, provided additional evidence of the accuracy of the temperature measurement, which is a critical aspect in obtaining accurate cross sections for reactions of metal atoms. The PL/GPID GIB instrument was used to continue the study of ion-molecule reactions that play a significant role in the low earth orbit spacecraft environment. One of the important reactions comprises collisions of atomic oxygen ions and water molecules. The extension of the study of these reactions to those with heavy water has been concluded.				
14. SUBJECT TERMS High temperature collision cell octopole Ion-metal atom charge transfer Atomic oxygen			15. NUMBER OF PAGES 12	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified		19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified
				20. LIMITATION OF ABSTRACT SAR

TABLE OF CONTENTS

1.0 ORIGINAL OBJECTIVES FOR YEAR 2	1
2.0 FORECAST FOR YEAR 2, AS PROPOSED AT THE END OF YEAR 1	1
3.0 SUMMARY OF PROGRESS	2
4.0 TECHNICAL ACTIVITIES	2
4.1 ION-METAL ATOM CHARGE TRANSFER-GIB	2
4.2 ION-MOLECULE REACTION STUDIES-GIB.....	3
4.3 CLUSTER ION STUDIES-GIB	4
4.4 NEW INSTRUMENT DEVELOPMENT.....	4
5.0 PATENT APPLICATION.....	5
6.0 CONFERENCES ATTENDED	5
7.0 PRESENTATIONS.....	5
8.0 PUBLICATIONS.....	6
9.0 TRAINING	6
10.0 FORECAST FOR YEAR 3.....	6

CONTROL OF CONTAMINANTS AROUND SPACECRAFT 1996 ANNUAL REPORT

The following report comprises a description of the activities performed in accordance with the contract between ORION International Technologies and the PL/GPID. Note that the Chemical Processes in the Space Environment Task of PL/WSSI, Spacecraft Interactions Branch, with which the contract was initiated, was transferred to PL/GPID during FY96.

1.0 ORIGINAL OBJECTIVES FOR YEAR 2

1. Complete the $O^+ + Ba$ experiments using the GIB apparatus.
2. Design and construct a water cluster ion source for the GIB instrument. Install and determine the optimal conditions for producing water cluster ions of the sizes of interest.
3. Continue development of the fast neutral source in the HTMS.
4. Continue the ongoing program for studying the hyperthermal ion-molecule reactions involved in the spacecraft-ionosphere interactions using the GIB machine.

2.0 FORECAST FOR YEAR 2, AS PROPOSED AT THE END OF YEAR 1

1. Assemble and test the high temperature octopole/collision cell apparatus. The initial testing will be done in the HTMS, if possible. This stage would involve testing of the cell and octopole heating elements, and assessing the temperature at various locations of the device. The high temperature octopole/collision cell operating characteristics will be tested after installation in the GIB instrument. Testing will take place after the completion of a series of GIB water cluster ion experiments currently being carried out by PL/WSSI personnel.
2. Use the high temperature octopole/collision cell apparatus to determine the $O^+ + Ba$ charge transfer cross section as a function of collision energy. This most important first stage may be followed up by attempting to elucidate the dynamics of the charge transfer process, through time-of-flight and emission studies.
3. Use the high temperature octopole/collision cell apparatus to study charge transfer reactions between important atmospheric ions, such as O^+ , N^+ , O_2^+ , NO^+ and N_2^+ , and other metals, such as Na and Mg, which have been observed in the ionosphere. In addition, the formation and chemistry of solvated metal clusters may be studied.
4. Use the GIB instrument, in its normal configuration, to continue the ongoing PL/WSSI program for studying hyperthermal ion-molecule reactions involved in the spacecraft-ionosphere interactions.

3.0 SUMMARY OF PROGRESS

1. The high temperature collision cell/octopole apparatus has been installed and calibrated. A manuscript, in which the apparatus is described and initial results are presented, has been submitted for publication in the Review of Scientific Instruments. An application for a patent of the high temperature apparatus has been initiated. The system is currently being used to study reactions of sodium with N_2^+ , O_2^+ , and NO^+ .
2. Knudsen Cell experiments were conducted in the HTMS to confirm the sodium vapor pressure curve and to determine the purity of the sodium metal sample intended for use in the ion-metal atom studies. The HTMS has since been dismantled, in accordance with changing priorities at PL/GPID.
3. The $O^+ + H_2O/D_2O$ study carried out in GIB instrument was extended, on the advice of reviewers, prior to publication of the work in Chemical Physics Letters.
4. A new cluster ion source has been installed in the GIB instrument and the first experiments, on the photodissociation and collision-induced dissociation of $(N_2O \cdot H_2O)^+$ cluster ions, have been completed.
5. Work has been initiated on the design of a new bakeable GIB instrument which will incorporate the features of the HTMS and will eventually house the HT8P. The new instrument will feature provisions for more flexibility in experiment capabilities and *in situ* sample purity determinations.

4.0 TECHNICAL ACTIVITIES

4.1 ION-METAL ATOM CHARGE TRANSFER-GIB

The high temperature octopole/collision cell apparatus (HT8P) was installed in the GIB instrument, along with the required cooling/power feedthroughs and a modified "injector lens". The radio frequency operating characteristics of the HT8P were optimized, and the collision cell effective interaction length was determined in a series of calibration experiments. The well known $Ar^+ + D_2 \rightarrow ArD^+ + D$ reaction was used in the calibration experiments, in which the HT8P was operated at temperatures up to 630 K (675°F). The collision cell gas density versus temperature dependence for measurements with a capacitance manometer was modeled theoretically and verified experimentally using the same reaction. This dependence, due to thermal transpiration, provided additional evidence of the accuracy of the temperature measurement, which is a critical aspect in obtaining accurate cross sections for reactions of metal atoms.

Evaluation of the performance of the new octopole/injector arrangement indicated an improvement in ion beam energy distribution from a typical energy width of 0.30 eV, in the old system, to a spread of as low

as 0.12 eV. Prior to installation of the HT8P, the operating characteristics of the octopole rod heaters were tested in the vacuum environment of the HTMS instrument. In addition, the temperature versus heating power dependence of the HT8P pole heaters was determined *in situ* by spot welding a thermocouple directly to one of the poles, and monitoring the temperature as all the poles were heated. The thermocouple was subsequently removed to allow normal operation of the HT8P.

Because of limitations on operating temperature due to the proximity of the HT8P to low temperature materials in the GIB instrument, the first metal to be studied using the HT8P was sodium which required operation at moderate temperatures (~460 K). Studies of the charge transfer reactions of sodium with N_2^+ , O_2^+ , and NO^+ were undertaken and are nearly completed. The new data represent a significant improvement over existing data and will make an important contribution to the effort to model the behavior of atmospheric/ionospheric metal species derived via meteor ablation.

During the sodium metal work, modifications to the GIB instrument to accommodate operation of the HT8P at higher temperatures were designed and the necessary hardware was fabricated. The modifications primarily involved replacing Teflon insulators with assemblies manufactured of stainless steel and MACOR or of alumina. In addition, the vacuum system was fitted with turbomolecular pumps.

The sodium metal studies provided an excellent test case for evaluating the use of atomic absorption spectrometry (AAS) as an independent method for determining the density of metal vapor in the collision cell. For sodium, AAS and the temperature/vapor pressure results were in good agreement, although better reproducibility was obtained with the latter method. AAS was found to be accurate and will be important in studies of metals for which vapor pressure data are not of high quality.

In support of the planned HT8P study of the $O^+ + Ba$ charge transfer cross section, additional experiments were done in the HTMS to improve our measurements of the Ba vapor pressure in the temperature range from 700 K to 900 K (427°C - 627°C). In addition, the measurement of barium vapor density using AAS was determined to be feasible, on the basis of the known spectroscopic characteristics of barium.

4.2 ION-MOLECULE REACTION STUDIES-GIB

The PL/GPID GIB instrument was used to continue the study of ion-molecule reactions that play a significant role in the low earth orbit spacecraft environment. One of the important reactions comprises collisions of atomic oxygen ions and water molecules. The extension of the study of these reactions to those with heavy water has been concluded. The isotope substitution was useful for elucidating aspects of the reaction dynamics for both the atom abstraction and charge transfer channels, and results from these experiments are included in publications 1 and 2, respectively, listed below. An attempt to extend the

isotopic substitution to include ^{18}O , on the advice of the reviewer for publication 1, failed to produce additional information.

4.3 CLUSTER ION STUDIES-GIB

The new cluster ion source, featuring a pulsed-supersonic nozzle crossed with a magnetically confined electron beam, was installed in the GIB instrument. Operation of the source was characterized, and optimal cluster formation conditions were determined. The system was first used to study $(\text{N}_2\text{O}\cdot\text{H}_2\text{O})^+$ cluster ions in photodissociation and collision-induced dissociation experiments. This system was chosen since it had been studied previously. The new photodissociation experiments utilized the new PL/GPID YAG-pumped OPO laser. The design and characterization of this system was performed by PL/GPID personnel; the experiments included participation by ORION personnel. A publication (No. 4, below) of the results of this study is in preparation.

4.4 NEW INSTRUMENT DEVELOPMENT

The HTMS apparatus was disassembled and design work begun for a new high temperature (bakeable to 525 K) instrument. The new instrument incorporates the essential elements of the old HTMS, i.e., sensitive detection of neutral species, and will house the HT8P apparatus for GIB studies. The existing GIB instrument will be refitted with the original room temperature octopole/collision cell apparatus. The essential elements of the design necessary to perform GIB work were completed, and some electronic and vacuum components were ordered/received. Specifically, the designs of the main chamber, the coupling of the magnetic sector instrument to the main chamber, and the injection lens system are completed. Furthermore, vacuum feedthroughs, stainless steel components and flanges, Wien velocity filter chamber and electronics, and an arbitrary waveform generator have been ordered. The remaining design work includes the quadrupole mass spectrometer detection system, the details of the feedthrough flange assemblies, and if funds are available, a new electron impact ionization source. The remaining equipment ordering, including voltage power supplies and gas handling equipment, will be completed in the next fiscal year, pending funding.

In addition, we have proposed a second application of the new instrument involving the development of a fast neutral beam source. This application builds on the expertise gained from detailed studies of the translational energy dependence and disposal of ion-molecule reactions. We proposed studying the translational and internal energy dependence of chemical reaction cross sections of fast neutral-molecule reactions relevant to hypersonic flows. An important aspect of the new experiment is the generation of intense, mass selected positive-ion beams with well-defined kinetic energy distributions at energies between 1 and 20 eV. The generation of intense positive-ion beams is an existing capability in our laboratory, where 10 nA ion beam currents at ~ 10 eV are routinely achieved. The ion beam is then

converted to a neutral beam in a cell containing a charge-transfer target gas. As the previous studies of our laboratory have shown, charge transfer can readily provide 10% neutralization with minimal energy and momentum transfer, thus producing a neutral beam with high directionality and well-defined translational energy.

In the case of N_2 production, our laboratory studies have shown that the state-to-state dynamics of $N_2^+ + H_2O$ charge-transfer collisions results in N_2 molecules that maintain the vibrational excitation of the parent ion. In addition, H_2O is an excellent charge-transfer target, because it can be efficiently pumped outside the cell, a necessity of this experiment. The neutral beam then passes through a second, thin cell containing a thermal target gas. The scattered and unscattered particles pass through an electron impact ionizer that is followed by efficient coupling of all ions to a quadrupole mass spectrometer for mass analysis and detection. Measurement of target gas density, primary neutral and secondary neutral intensities yield absolute cross sections. The design work to add these additional features is currently underway and will continue, pending funding for FY97 and PL/GPID priorities.

5.0 PATENT APPLICATION

A patent application entitled "High-Temperature Octopole Ion Guide with Coaxially Heated Rods", by Dale J. Levandier & Rainer A. Dressler, is currently awaiting processing at the Air Force ESC/JAZ office.

6.0 CONFERENCES ATTENDED

1. 1996 Air Force Office of Scientific Research Contractors' Review, Boulder, CO, 96 JUN 02 - 96 JUN 05.
2. The Gordon Research Conference on Atomic and Molecular Interactions, New London, NH, 96 JUN 30 - 96 JUL 05
3. The 14th International Symposium on Gas Kinetics, Leeds, UK, 96 SEP 07 - 96 SEP 12.

7.0 PRESENTATIONS

1. "High Temperature Guided-Ion Beam Experiment for Ion-Metal Vapor Collision Studies", Dale J. Levandier, Skip Williams, Rainer A. Dressler & Edmond Murad - a poster presented by Dale Levandier at the 1996 AFOSR Contractors' Review.
2. "Collision-Induced Dissociation and Photodissociation Studies of the $(N_2O \cdot H_2O)^+$ Cluster Ion", Michael J. Bastian, Rainer A. Dressler, Dale J. Levandier & Edmond Murad - a poster presented by Dale Levandier at the 14th International Symposium on Gas Kinetics.

8.0 PUBLICATIONS

1. A study of the isotope effects in the reaction $O^+ + H_2O/D_2O \rightarrow OH^+/OD^+ + OH/OD$ using guided-ion beams.
Dale J. Levandier, Rainer A. Dressler & Edmond Murad
Chemical Physics Letters, Vol. 251, pp. 174-181, 1996.
2. Empirical model of the state-to-state dynamics in near-resonant hyperthermal $X^+ + H_2O$ charge-transfer reactions.
Rainer A. Dressler, Michael J. Bastian, Dale J. Levandier & Edmond Murad
International Journal of Mass Spectrometry and Ion Processes, accepted for publication.
3. A high-temperature octopole ion guide for measuring absolute cross sections of ion-metal atom reactions.
Dale J. Levandier, Rainer A. Dressler & Edmond Murad
Review of Scientific Instruments, submitted for publication, 96 SEP.
4. Low energy collision-induced dissociation and photodissociation studies of the $(N_2O \cdot H_2O)^+$ cluster ion.
Michael J. Bastian, Rainer A. Dressler, Dale J. Levandier, Edmond Murad, Felician Muntean & Peter B. Armentrout
In preparation.

9.0 TRAINING

1. 96 MAY. Hazardous Waste Management Training - Refresher pursuant to EPA 40 CFR 260-279 (DL). The 4 hour course was conducted at Hanscom AFB by Institute for Environmental Education, Inc.
2. 96 MAY. Hazardous Waste Management Training - Initial pursuant to EPA 40 CFR 260-279 (SW). The 7 hour course was conducted at Hanscom AFB by Institute for Environmental Education, Inc.

10.0 FORECAST FOR YEAR 3

1. Conclude the HT8P-GIB study of sodium atom reactions.
2. Use the HT8P to determine the $O^+ + Ba$ charge transfer cross section as a function of collision energy.
3. Use the HT8P to study charge transfer reactions between important atmospheric/ionospheric ions, such as O^+ , N^+ , O_2^+ , NO^+ and N_2^+ , and other metals, such as Mg and Ca, which have been observed in the ionosphere. In addition, the formation and chemistry of solvated metal cluster ions may be

studied. These species are expected to play important roles in metal transport phenomena in the upper atmosphere/ionosphere.

4. Use the GIB instrument, in its normal configuration or with the cluster ion source, to continue the ongoing PL/GPID program for studying hyperthermal ion-molecule reactions involved in the spacecraft-ionosphere interactions. These experiments will include photodissociation and collision-induced dissociation studies of excited molecular ion moieties of the sort generated in spacecraft thruster firings, and produced for the experiments in the supersonic nozzle cluster ion source. As a test case, we have proposed generating acetylene-water ion clusters in the cluster ion source. In the source region this moiety may undergo reactions producing isomeric structures, namely aldehyde and enol species, that can be distinguished in CID experiments.
5. Continue the development of the new GIB apparatus, including the design work on the fast neutral beam source.